

Population genetic characteristic of horses of Mugalzhar breed by STR-markers.

Seleuova LA*, Naimanov DK, Jaworski Z, Aubakirov MZH, Mustafin MK, Mustafin BM, Safronova OS, Baktybaev GT, Turabaev AT, Domatski VN

Baitursynov State University, Kostanay, Baytursynov Str., 47, Kazakhstan

Abstract

One of the most important tasks of improvement of horse breeds is investigation into genetic determinants responsible for formation of high productivity and use of genetic monitoring upon selection control. Nowadays genome selection is the most advanced estimation method of breeding abilities. This article presents results of genetic testing of DNA of Mugalzhar horses by 16 microsatellite loci, from 5 to 13 alleles have been identified. Genetic diversity (N_v) of modern population of purebred brood horses was at the level of 7.882, polymorphism (A_e) equalled to 4.457, heterozygosity (H_o)-to 0.754, H_e -to 0.775, fixation index (F_{is})-to 0.027; it referred to a cluster of native/local breeds. Polymorphism varied from 2.785 to 7.442; actually observed heterozygosity from 0.138 to 0.931; theoretically expected one-from 0.641 to 0.866. The number of active alleles in population (N_a) was equal to 134.0.

Keywords: Genome selection, Horse breed, Molecular genetic markers, Polymorphism, Population, Primer.

Accepted on September 26, 2018

Introduction

Monitoring of population genetic polymorphism is one of the most important tasks of preservation and reproduction of various agricultural breeds including horses [1,2]. In this regard characteristics of population genetic structure can be the first step to breed preservation and reproduction, in addition it improves cattle breeding [3,4].

Such analysis is carried out on using molecular genetic technologies based on various DNA markers [3,5]. This approach expands significantly capabilities of population genetic analysis, makes it possible to reveal cross- and inbreed variety of separate genome segments, presents breed genetic structure and serves as excellent tool for investigation into genetic variance in the frames of genus or species [5-10].

In 1998 year (Order No. 156 of Ministry of Agriculture of Kazakhstan dated December 30, 1998) the researchers and horse breeders of Kazakhstan, using pure breeding and improvement of Jabe horse breed, created specialized dual-purpose Mugalzhar horse breed which could provide high quality meat and kumis during yearlong grassland farming [11,12].

Taking into account high value of the domestic Mugalzhar horse breed and increased demand for meat and dairy products of horse breeding, it would be reasonable to preserve and to improve the Mugalzhar breed in Kazakhstan on a wider range, this stipulated the targets of our work.

Practical value of the studies: the obtained results evidence activity of genetic processes upon horse breeding in «Sholak Espe» farm, high polymorphism and high genetic diversity of the considered horse breed. The available data on horse genotypes make it possible to perform more efficient selection aimed at preservation of genetic peculiarities of lines (family groups) and support of inbreed differentiation.

Materials and Methods

The studies were performed with samples of biological material of 70 Mugalzhar horses bred in «Sholak Espe» farm in Karaganda region (Central Kazakhstan). We used hair bulbs of the tested horses as biological material for DNA isolation.

These studies were performed in test laboratory of TOO «Kazak Tulpary» (Ministry of agriculture of Kazakhstan, Kostanay) certified according to ISO/IEC 17025-2009.

Genome DNA was isolated from hair bulbs of horses with proper modifications using reagents produced by OOO «Izogen», Russia, according to the producer recommendations. Genetic typing of the horses was carried out using a set of 16 microsatellite loci recommended by International Society for Animal Genetics (ISAG): *HVL20*, *HTG4*, *AHT4*, *HMS7*, *HTG6*, *AHT5*, *HMS6*, *ASB23*, *ASB2*, *HTG10*, *HTG7*, *HMS3*, *HMS2*, *ASB17*, *LEX3*, *HMS1*. The amplification products were identified using an ABI Prism 310 (Applied Biosystems, USA) genetic analyzer on the basis of capillary electrophoresis and laser detection. The acquired graphical results were processed

using Gene Mapper 4.0 software. Polymorphism was characterized by the following indices: allele frequency of the considered loci calculated by maximum likelihood formula ($p_A \rightarrow$ Equation (1)) and genotype frequency (PAA \rightarrow Equation (2)), actually observed ($H_o \rightarrow$ Equation (3)) and theoretically expected heterozygosity ($H_e \rightarrow$ Equation (4)) with consideration for the Hardy-Weinberg law H_o -according to Ney, H_e (with C_a according to Robertson \rightarrow Equation (5)), as well as average heterozygosity for populations, fixation index (Fis \rightarrow Equation (6)), polymorphism level (Ae \rightarrow Equation (7)), average number of alleles in locus ($N_v \rightarrow$ Equation (8)).

$$p_A = (2n_{AA} + n_{AB} + n_{AC}) / 2N \rightarrow (1)$$

where $2n_{AA}$ was the double number of homozygotes; n_{AB} , n_{AC} were the number of heterozygotes; $2N$ was the double number of analysed animals in the selection [13].

$$p_{AA} = n_{AA} / N \rightarrow (2)$$

where p_{AA} was the y frequency of genotypes of the considered DNA microsatellite loci; n_{AA} was the number of animals with genotype AA [13].

$$H_o = h_j / n \rightarrow (3)$$

where H_o was the actually observed heterozygosity for one locus; h_j was the number of heterozygote genotypes in locus; n was the total number of genotypes in locus [14].

$$H_e = 1 - C_a \rightarrow (4)$$

Table 1. Polymorphism of the considered microsatellite loci of DNA of Mugalzhari horses.

Microsatellite locus	Expected homozygosity (C_a)	Polymorphism (Ae)	Observed heterozygosity (H_o)	Expected heterozygosity (H_e)	Fixation index (Fis)	Number alleles (N_a)	of
HVL20	0.182	5.497	0.931	0.818	-0.138	11	
HTG4	0.308	3.247	0.724	0.692	-0.046	6	
AHT4	0.162	6.184	0.862	0.838	-0.028	8	
HMS7	0.303	3.298	0.724	0.697	-0.039	7	
HTG6	0.260	3.849	0.690	0.740	0.068	6	
AHT5	0.235	4.247	0.690	0.765	0.098	6	
HMS6	0.359	2.785	0.724	0.641	-0.130	7	
ASB23	0.137	7.313	0.931	0.863	-0.079	11	
ASB2	0.160	6.253	0.724	0.840	0.138	10	
HTG10	0.184	5.426	0.828	0.816	-0.015	10	
HTG7	0.249	4.014	0.690	0.751	0.082	5	
HMS3	0.189	5.289	0.828	0.811	-0.021	7	
HMS2	0.170	5.881	0.897	0.830	-0.080	8	
ASB17	0.134	7.442	0.828	0.866	0.044	13	
LEX3	0.240	4.163	0.138	0.760	0.818	7	
HMS1	0.293	3.412	0.897	0.707	-0.268	5	
Average	0.2243	4.4574	0.7546	0.7757	0.0272	134.0	

where H_e was the expected heterozygosity; C_a was the expected homozygosity, it was determined on the basis of coefficient of homozygosity by the Robertson formula [15].

$$C_a = \sum np^2 \rightarrow (5)$$

where C_a was the expected homozygosity; p was the gene frequency of alleles; n was the number of alleles in locus [15].

$$F_{is} = 1 - (H_o / H_e) \rightarrow (6)$$

where H_o was the actually observed heterozygosity; H_e was the expected heterozygosity [15].

$$A_e = 1 / (\sum p_{ij}^2) \rightarrow (7)$$

where A_e was the level of polymorphism (active effective alleles); p was the occurrence frequency of the j^{th} allele for locus I, and summation was applied for n alleles [14].

$$N_a = 1 / C_a \rightarrow (8)$$

where N_a was the number of active alleles in population (polymorphism); C_a was the level of expected homozygosity of horses by polymorphic alleles [13].

Results and Discussion

While analyzing the allele pool of the considered horses by 16 DNA microsatellites, we obtained data characterizing polymorphism of each marker (Table 1).

On the basis of the considered 16 STR-loci from 5 to 13 alleles were identified. According to Table 1, it was established that average polymorphism of locus (Ae) was 4.457, hence, all loci were subdivided into two groups. The first group was comprised of loci with the level of polymorphism below average: *HMS6*, *HTG4*, *HMS7*, *HMS1*, *HTG6*, *HTG7*, *LEX3*, *AHT5*. Minimum value was that of *HMS6*: 2.785. The second group was comprised of loci with the level of polymorphism above average: *ASB17*, *ASB23*, *ASB2*, *AHT4*, *HMS2*, *HVL20*, *HTG10*, *HMS3*. Maximum polymorphism was that of *ASB17*: 7.442.

In order to obtain precise estimation of population variance, the index of expected heterozygosity was introduced which considered the level of allele diversity. In this regard we estimated the actually observed and the theoretically expected heterozygosity calculated by 16 loci, the respective average indices were 0.7546 and 0.7757.

The highest actually observed heterozygosity (Ho) was that of locus *HVL20* (0.931), and that of theoretically expected heterozygosity (He)-locus *ASB17* (0.866), whereas the minimum heterozygosity was that of locus *LEX3* (0.138), located on sex chromosome, and the minimum expected heterozygosity was that of locus *HMS6* (0.641).

The calculated fixation index *Fis* made it possible to establish interrelation among the considered horses of Mugalzhar breed of Kozhamberdy selection. We revealed that the deficiency of heterozygotes was observed for loci *HTG6*, *AHT5*, *ASB2*, *HTG7*, *ASB17*, *LEX3* (Figure 1).




Figure 1. Expected (He), observed (Ho) heterozygosity and fixation index (Fis) in population of Mugalzhar horses of Kozhamberdy selection.

For preservation of genetic inbred diversity, average number of alleles (Nv) for all the considered markers in the breed were of great concern. In the Mugalzhar breed it was 7.882.

In addition, we counted the number of active alleles in population by Na, which also characterized the level of polymorphism. This variable was inversely proportional to the Robertson homozygosity coefficient. The highest level of polymorphism was observed for the Mugalzhar horses on locus *ASB17* [13]. This variable in the aggregate for all considered loci amounted to 134.0.

While studying published information, we revealed that the method of determination of genetic polymorphism of horses and other farm livestock using marker microsatellites was widely applied in practice. Thus, Chasymova et al. while estimating genetic variation in populations of Tuvinian horses, detected from 4 to 9 alleles in the considered microsatellite loci, in our study this parameter was from 5 to 13 alleles. Microsatellite loci *ASB17*, *AHT4*, *VHL20*, *ASB2* and *ASB23* were presented by the most varied range of alleles which had been confirmed by our results. However, in Mugalzhar horses the loci *HMS2*, *HMS3* were also marked as loci of high polymorphism [16].

Mahrous et al. reported about high information content of microsatellite loci *AHT4*, *HTG10*, *ABS2*, *ABS23* which also corresponded with our experimental results [17].

It is possible to state that application of microsatellite markers has high resolution ability making it possible to estimate consistency of the considered stock, to perform more efficient selection of animals, to control inheritance of valuable properties.

Conclusion

On the basis of the considered 16 STR-loci from 5 to 13 alleles were identified. The average polymorphism of locus (Ae) was 4.457, hence, all loci were subdivided into two groups. The first group was comprised of loci with the level of polymorphism below average: *HMS6*, *HTG4*, *HMS7*, *HMS1*, *HTG6*, *HTG7*, *LEX3*, *AHT5*. Minimum value was that of *HMS6*: 2.785. The second group was comprised of loci with the level of polymorphism above average: *ASB17*, *ASB23*, *ASB2*, *AHT4*, *HMS2*, *HVL20*, *HTG10*, *HMS3*. Maximum polymorphism was that of *ASB17*: 7.442. The actually observed heterozygosity was from 0.138 to 0.931; the theoretically expected one was from 0.641 to 0.866, average indices were 0.7546 and 0.7757, respectively.

The highest actually observed heterozygosity (Ho) was that of locus *HVL20* (0.931), and that of theoretically expected heterozygosity (He)-locus *ASB17* (0.866), whereas the minimum heterozygosity was that of locus *LEX3* (0.138), located on sex chromosome, and the minimum expected heterozygosity was that of locus *HMS6* (0.641).

The calculated fixation index *Fis* made it possible to establish interrelation among the considered horses of Mugalzhar breed of Kozhamberdy selection. We revealed that the deficiency of heterozygotes was observed for loci *HTG6*, *AHT5*, *ASB2*, *HTG7*, *ASB17*, *LEX3*. Average number of alleles (Nv) for all considered markers in Mugalzhar horse breed was 7.882.

In addition, we counted the number of active alleles in population by Na, which also characterized the level of polymorphism. This variable was inversely proportional to the Robertson homozygosity coefficient. The highest level of

polymorphism was observed for the Mugalzhars on locus *ASB17* [13]. This variable in the aggregate for all considered loci amounted to 134.0.

References

1. Barcaccia G, Felicetti M, Galla G, Capomaccio S, Supplizi AV. Molecular analysis of genetic diversity, population structure and inbreeding level of the Italian Lipizzan horse. *Livestock Science* 2013; 151: 124-133.
2. Metlitska OI. The Methodology of DNA gene pools certification of farm animals for hypervariable loci in the genome. Poltava 2012.
3. Kalashnikov VV, Zaitsev AM, Khrabrova LA. Genetic identification and origin monitoring in horse breeding. Scientific support of development and efficiency of brood, sports and productive horse breeding in Russia and CIS countries: Proceedings of International Conference devoted to the 75th anniversary of Prof. Koveshnikov, 2014.
4. Suprun IA. Evaluation of phylogenetic connections in livestock breeding. Golovna 2014.
5. Rothschild MF, Plastow GS. Applications of genomics to improve livestock in the developing world. *Livestock Science* 2014; 166: 76-83.
6. Ganopoulos I, Kalivas A, Kavroulakis N, Xanthopoulou A, Mastrogianni A, Koubouris G, Madesis P. Genetic diversity of Barbary fig (*Opuntia ficus-indica*) collection in Greece with ISSR molecular markers. *Plant Gene* 2015; 2: 29.
7. Korbin M, Kuras A, Zurawicz E. Fruit plant germplasm characterisation using molecular markers generated in RAPD and ISSR-PCR. *Cell Mol Biol Lett* 2002; 7: 785-794.
8. Sarla N, Bobba S, Siddiq EA. ISSR and SSR markers based on AG and GA repeats delineate geographically diverse *Oryza nivara* accessions and reveal rare alleles. *India Curr Sci* 2003; 84: 683.
9. Stolpovskiy YA. Application inter-SSR DNA analysis to assess population structure, identification and similarity of gene pools of species and breeds of domesticated animals. *Genetics* 2010; 46: 825-833.
10. Zabek T, Nogay A, Radko A, Nogay J, Slota E. Genetic variation of Polish endangered Bilgoraj horses and two common horse breeds in microsatellite loci. *Appl Gene* 2005; 46: 299-305.
11. Satybaldin AA. Current state of horse breeding and horse sports in Kazakhstan. The First International Conference Kostanay 2002.
12. Turabaev A. New lines in Kulandy interbreed type of Mugalzhars horse breed. *Novosti Nauki Kazakhstana* 2011; 156.
13. Estimation procedure of genetic diversity and genetic similarity of farm and local breeds. Divovo 2011.
14. Kantanen J. Genetic diversity and population structure of 20 north European cattle breeds. *J Heredity* 2000; 91: 446-457.
15. Methodology of application of DNA analysis of horses for estimation of genetic resources in horse breeding. Divovo 2011.
16. Chasymova RB, Khrabrova LA, Zaitsev AM, Makarova EYu, Fedorov YuN, Ludu BM. Estimation of genetic variety in populations of Tuvian horses according to loci of blood system and microsatellite DNA. *Sel'skokhozyaistvennaya Biologiya* 2017; 52: 679-685.
17. Mahrous KF, Hassanane M, Mordy A, Shafey HI, Hassan N. Genetic variations in horse using microsatellite markers. *J Genetic Eng Biotechnol* 2011; 9: 103-109.

*Correspondence to

Seleuova LA
Baitursynov State University
Kostanay
Kazakhstan